arXiv:1501.02547v2 [math.AT] 29 Jun 2016

THE HOCHSCHILD HOMOLOGY OF $A(1)$.

ANDREW SALCH

Abstract. We compute the Hochschild homology of $A(1)$, the subalgebra of the 2-primary Steenrod algebra generated by the first two Steenrod squares, $\mathrm{Sq}^1, \mathrm{Sq}^2$. The computation is accomplished using several May-type spectral sequences.

Contents

1. Introduction. 1
2. Construction of May-type spectral sequences for Hochschild homology. 4
3. The May and abelianizing filtrations. 9
4. Running the HH-May and abelianizing spectral sequences. 11
4.1. Input. 11
4.2. d_1-differentials. 14
4.3. d_2-differentials. 16
References 19

1. Introduction.

The 2-primary Steenrod algebra A, that is, the algebra of stable natural endomorphisms of the mod 2 cohomology functor on topological spaces, has generators $\mathrm{Sq}^1, \mathrm{Sq}^2, \mathrm{Sq}^3, \ldots$, the *Steenrod squares*. The subalgebra of A generated by the first two Steenrod squares, Sq^1 and Sq^2, is called $A(1)$, and $A(1)$ is an eight-dimensional, graded, noncommutative (not even graded-commutative), co-commutative Hopf algebra over $\mathbb{F}_2$. The homological algebra of $A(1)$-modules effectively determines, via the Adams spectral sequence, the 2-complete homotopy theory of spaces and spectra smashed with the connective real K-theory spectrum ko. These ideas are all classical; an excellent reference for the Steenrod algebra is Steenrod's book [12], and an excellent reference for $A(1)$-modules and the Adams spectral sequence is the third chapter of Ravenel's book [11].

As a student of homotopy theory, when I first learned the definition of Hochschild homology of algebras, my first reaction was to try to compute the Hochschild homology of $A(1)$. I know at least three other homotopy theorists who have told me that they had the same reaction when learning about Hochschild homology! Computing $HH_*(A(1), A(1))$, however, is a nontrivial task, and it seems that this computation has never been successfully done[1].

Date: January 2015.

[1]Bökstedt, in his extremely influential unpublished paper on topological Hochschild homology, computes the Hochschild homology of $\pi_*(H\mathbb{F}_p \wedge H\mathbb{F}_p)$, i.e., the Hochschild homology of *the linear dual* of the entire Steenrod algebra, but this is very straightforward, since the dual of the Steenrod algebra is polynomial at $p = 2$ and polynomial tensored with exterior at $p > 2$. For the same reason, it is also easy to compute the Hochschild

In this paper we compute $HH_*(A(1), A(1))$ by using two different filtrations on $A(1)$ and studying the spectral sequences in Hochschild homology arising from these filtrations. These spectral sequences are the analogues in Hochschild homology of J. P. May's spectral sequence for computing Ext over the Steenrod algebra (see [7]), so we think of these as "May-type" spectral sequences.

The problem of computing $HH_*(A(1), A(1))$ is made rather difficult by the fact that $A(1)$ is noncommutative and so $HH_*(A(1), A(1))$ does not inherit a product from the shuffle product on the cyclic bar complex, and as a consequence, the May-type spectral sequence converging to $HH_*(A(1), A(1))$ that one would construct in the most naïve way is not *multiplicative*, i.e., it does not have a product satisfying a Leibniz rule. This makes the computation of differentials in that spectral sequence basically intractable. Instead, we take the linear dual of the standard Hochschild chain complex on $A(1)$, and we use the cocommutative coproduct on $A(1)$ to give the cohomology of this linear dual cochain complex a product structure arising from the coproduct on $A(1)$ and the linear dual of the Alexander-Whitney map. In Proposition 4.3 we set up *multiplicative* spectral sequences computing the cohomology of the linear dual cochain complex of the standard Hochschild chain complex of $A(1)$. By an easy universal coefficient theorem argument (Proposition 2.5), this cohomology is the $\mathbb{F}_2$-linear dual of the desired Hochschild homology $HH_*(A(1), A(1))$.

We then compute the differentials in these spectral sequences. In the end there are nonzero d_1 and d_2 differentials, and no nonzero differentials on any later terms of the spectral sequences. In 4.2.1 and 4.3.1 we present charts of the E_2 and $E_3 \cong E_\infty$-pages of the the relevant spectral sequences. Our charts are drawn using the usual Adams spectral sequence conventions, described below. This is the most convenient format if, for example, one wants to use this Hochschild homology as the input for an Adams spectral sequence, and it also makes it easier to see the natural map from this Hochschild homology to the classical Adams spectral sequence computing $\pi_*(ko)^\wedge_2$, the 2-complete homotopy groups of the connective real K-theory spectrum ko, in Proposition 4.4 and in the charts 4.3.1 and 4.3.2.

In particular, the chart 4.3.1 is a chart of the ($\mathbb{F}_2$-linear dual of the) Hochschild homology of $A(1)$, and gives our most detailed description of $HH_*(A(1), A(1))$. We reproduce that chart here:



homology of *the linear dual* of $A(1)$. But the linear dual of $A(1)$ is a completely different ring than $A(1)$, so this sheds no light on the Hochschild homology of $A(1)$ itself!

The vertical axis is homological degree, so the row s rows above the bottom of the chart is the associated graded $\mathbb{F}_2$-vector space of a filtration on $HH_s(A(1), A(1))$. The horizontal axis is, following the tradition in homotopy theory, the Adams degree, i.e., the topological degree (coming from the topological grading on $A(1)$) minus the homological degree. The horizontal lines in the chart describe comultiplications by certain elements in the linear dual Hopf algebra $\hom_{\mathbb{F}_2}(A(1), \mathbb{F}_2)$ of $A(1)$, and the nonhorizontal lines describe certain operations in the linear dual of $HH_*(A(1), A(1))$, described in Convention 4.6. The entire pattern described by this chart is repeated every four vertical degrees and every eight horizontal degrees: there is a periodicity class (not pictured) in bidegree $(4, 8)$.

Information about the $\mathbb{F}_2$-vector space dimension of $HH_*(A(1), A(1))$ in each grading degree is provided by Theorem 4.10, which we reproduce below (we do not describe any ring structure on $HH_*(A(1), A(1))$ because $A(1)$ is noncommutative and so there is no ring structure on its Hochschild homology induced by shuffle product on cyclic chains):

Theorem. *The $\mathbb{F}_2$-vector space dimension of $HH_n(A(1), A(1))$ is:*

$$\dim_{\mathbb{F}_2} HH_n(A(1), A(1)) = \begin{cases} 2n+5 & \text{if } 2 \mid n \\ 2n+6 & \text{if } n \equiv 1 \mod 4 \\ 2n+4 & \text{if } n \equiv 3 \mod 4. \end{cases}$$

Hence the Poincaré series of the graded $\mathbb{F}_2$-vector space $HH_*(A(1), A(1))$ is

$$\frac{5 + 8s + 9s^2 + 10s^3 + \frac{8s^4}{1-s}}{1 - s^4}.$$

If we additionally keep track of the extra grading on $HH_*(A(1), A(1))$ coming from the topological grading on $A(1)$, then the Poincaré series of the bigraded $\mathbb{F}_2$-vector space $HH_{*,*}(A(1), A(1))$ is

$$\left((1+u)\left(1 + u^2 + su(1+u^2+u^5) + s^2u^2(1+2u^5+u^7) + s^3u^3(1+u^4+u^5+u^6+u^9) \right.\right.$$
$$\left.\left. + \frac{s^4u^4(1+u^4+u^5+u^9)}{1-su}\right) + u^6 + su^2 + su^8 + s^2u^4\right)\frac{1}{1-s^4u^{12}}$$

where s indexes the homological grading and u indexes the topological grading.

Our computation of $HH_*(A(1), A(1))$ can be used as the input for other spectral sequences in order to make further computations. For example, one could use it as input for the Connes spectral sequence, as in 9.8.6 of [13], computing the cyclic homology $HC_*(A(1), A(1))$. This is probably of limited utility, however, since $A(1)$ is an algebra over a field of characteristic 2, so the cyclic homology of $A(1)$ is probably not a good approximation to the algebraic K-theory of $A(1)$. Instead one ought to compute the topological cyclic homology of $A(1)$. For this, one could use our computation of $HH_*(A(1), A(1))$ as input for the Pirashvili-Waldhausen spectral sequence, as in [10], computing the topological Hochschild homology $THH_*(A(1), A(1))$, and then one would need to run the necessary homotopy fixed-point spectral sequences to compute $\pi_*(THH(A(1), A(1))^{C_{p^n}})$ and then $TR_*(A(1))$ and finally $TC_*(A(1))$, which, using McCarthy's theorem (see [3]), gives the 2-complete algebraic K-groups $K_*(A(1))^\wedge_2$ (the algebraic K-groups completed away from 2 ought to be much easier: an appropriate noncommutative version of Gabber rigidity, if true, would imply that they vanish in positive degrees. See IV.2.10 of [14]). See e.g. [5] for a survey of trace method computations of this kind. Those computations are entirely outside the scope of the present paper.

We remark that our methods also admit basically obvious extensions to methods for computing $HH_*(A(n), A(n))$ for arbitrary n, but one sees that for $n > 1$, carrying out such computations would be a daunting task. Our HH-May spectral sequence of Proposition 4.3 surjects on to the classical May spectral sequence computing $\mathrm{Ext}^*_{A(1)}(\mathbb{F}_2, \mathbb{F}_2)$, and for the same reasons, the $n > 1$ analogue of our HH-May spectral sequence maps naturally to the classical May spectral sequence computing $\mathrm{Ext}^*_{A(n)}(\mathbb{F}_2, \mathbb{F}_2)$. We suspect that this map is still surjective for $n > 1$, although we have made no attempt to verify this. Consequently the computation of $HH_*(A(n), A(n))$ using our methods is of at least the same level of difficulty as the computation of $\mathrm{Ext}^*_{A(n)}(\mathbb{F}_2, \mathbb{F}_2)$. For $n = 2$ this is already quite nontrivial.

It is worth pointing out that, in [2], J.-L. Brylinski constructs a spectral sequence related to the present paper's abelianizing spectral sequence. Brylinski's spectral sequence computes the Hochschild homology of a noncommutative algebra over a field of characteristic zero, for the purposes of studying Poisson manifolds, and remarks that "[e]xamples show that this spectral sequence tends to degenerate at E^2," in particular, that Brylinski's spectral sequence collapses at the E^2-term for the algebra of differential operators on an algebraic or complex-analytic manifold. In the present paper's computation of $HH_*(A(1), A(1))$ we instead get collapse one term later, at E_3 rather than E_2.

Remark 1.1. Given a field k and a graded k-algebra A, there are *two* notions of the Hochschild homology $HH_*(A, A)$ which are in common circulation:

- one can forget the grading, and simply consider the Hochschild homology of the underlying ungraded k-algebra of A. This is the right thing to do in many applications of Hochschild homology in classical algebra; for example, if one wants to use trace methods to compute the algebraic K-groups of (the underlying ungraded algebra of) A. (The computations in [4] are an excellent example of this.)
- Alternatively, one can instead compute the "graded algebra Hochschild homology," which incorporates a sign convention into the cyclic bar complex. This is the right thing to do in many applications of Hochschild homology in algebraic topology; for example, if one wants to use Bökstedt's spectral sequence to compute topological Hochschild homology of a ring spectrum. (The computations in [8] are an excellent example of this.)

Since $A(1)$ has characteristic 2, sign conventions are irrelevant, and both notions of Hochschild homology coincide. This makes the computations in this paper equally applicable in classical algebra as in algebraic topology. Any future odd-primary analogues of these computations, however, would require that one carry out "ungraded" HH_* computations separately from the "graded" HH_*-computations.

2. Construction of May-type spectral sequences for Hochschild homology.

Proposition 2.1. (May spectral sequence for Hochschild homology.) *Let k be a field, A an algebra, and*

$$F^0A \supseteq F^1A \supseteq F^2A \supseteq \dots \tag{2.0.1}$$

a filtration of A which is multiplicative, that is, if $x \in F^mA$ and $y \in F^nA$, then $xy \in F^{m+n}A$. Then there exists a spectral sequence

$$E_1^{s,t} \cong HH_{s,t}(E_0A, E_0A) \Rightarrow HH_s(A, A)$$
$$d_r^{s,t} : E_r^{s,t} \to E_r^{s-1,t+r}.$$

The bigrading subscripts $HH_{s,t}$ are as follows: s is the usual homological degree, while t is the May degree, defined and computed as follows: given a homology class $x \in HH_s(E^0A, E^0A)$,

its May degree is the total degree (in the grading on E^0A induced by the filtration on A) of any homogeneous cycle representative for x in the standard Hochschild chain complex.

This spectral sequence enjoys the following additional properties:

- *If the filtration 2.0.1 is finite, i.e., $F^nA = 0$ for some $n \in \mathbb{N}$, then the spectral sequence converges strongly.*
- *If A is also a graded k-algebra and the filtration layers F^nA are generated (as two-sided ideals) by homogeneous elements, then this spectral sequence is a spectral sequence of graded k-vector spaces, i.e., the differential preserves the grading.*
- *If A is commutative, then so is E_0A, and the input for this spectral sequence has a ring structure given by the usual shuffle product on the Hochschild homology of a commutative ring (see e.g.* [13]*), and the spectral sequence is multiplicative, i.e., the differentials in the spectral sequence obey the graded Leibniz rule. Furthermore, the product in the spectral sequence converges to the product induced on the associated graded by the usual shuffle product on $HH_*(A, A)$ (this is the usual situation in spectral sequences of differential graded algebras).*
- *The differential in the spectral sequence is (like any other spectral sequence of a filtered chain complex) computed on a class $x \in HH_{*,*}(E_0A, E_0A)$ by computing a homogeneous cycle representative y for x in the standard Hochschild chain complex for E_0A, lifting y to a homogeneous chain $\tilde{y}$ in the standard Hochschild chain complex for A, applying the Hochschild differential d to $\tilde{y}$, then taking the image of $d\tilde{y}$ in the standard Hochschild chain complex for E_0A.*

Proof. Let $CH_\bullet(A, A)$ denote the standard Hochschild chain complex of A, and let $F^nCH_\bullet(A, A)$ denote the sub-chain-complex of $CH_\bullet(A, A)$ consisting of all chains of total filtration degree $\leq n$. Our May spectral sequence is now simply the spectral sequence of the filtered chain complex

$$CH_\bullet(A, A) = F^0CH_\bullet(A, A) \supseteq F^1CH_\bullet(A, A) \supseteq F^2CH_\bullet(A, A) \supseteq \dots. \tag{2.0.2}$$

If A is commutative, then filtration 2.0.1 being multiplicative implies that filtration 2.0.2 is a multiplicative filtration of the differential graded algebra $CH_\bullet(A, A)$, with product given by the shuffle product. It is standard (see. e.g. [6]) that the spectral sequence of a multiplicatively-filtered DGA is multiplicative.

The product on the spectral sequence being given by the shuffle product is due to the naturality of the construction of $CH_\bullet(A, A)$ in the choice of k-algebra A: if A is commutative, then the multiplication map $A \otimes_k A \to A$ is a morphism of k-algebras, hence we get a map of chain complexes

$$CH_\bullet(A \otimes_k A, A \otimes_k A) \to CH_\bullet(A, A),$$

which we compose with the Eilenberg-Zilber (i.e., "shuffle") isomorphism

$$CH_\bullet(A, A) \otimes_k CH_\bullet(A, A) \xrightarrow{\cong} CH_\bullet(A \otimes_k A, A \otimes_k A).$$

The other claims are also all standard about a spectral sequence of a filtered chain complex, except perhaps the strong convergence claim, which we easily prove as follows: suppose the filtration 2.0.1 satisfies $F^nA = 0$ for some $n \in \mathbb{N}$. Then the group of i-cycles $CH_i(A, A) \cong A^{\otimes_k i+1}$ has no nonzero elements of filtration greater than $(n + 1)i$. So the filtration in E_∞ of $HH_i(A, A)$ is a finite filtration, and the column in the spectral sequence converging to $HH_i(A, A)$ is zero after the $E_{(n+1)i+2}$-page. So the spectral sequence converges strongly. □

Definition 2.2. *Let k be a field and A a coalgebra over k with comultiplication map $\Delta : A \to A \otimes_k A$. By the* cyclic cobar construction *on A we mean the cosimplicial k-vector space*

$$A \rightrightarrows A \otimes_k A \;\substack{\rightarrow\\\rightarrow\\\rightarrow}\; A \otimes_k A \otimes_k A \;\substack{\rightarrow\\\rightarrow\\\rightarrow\\\rightarrow}\; \dots$$

with coface maps $d^0, d^1, \dots, d^n : A^{\otimes_k n} \to A^{\otimes_k n+1}$ given by

$$d^i(a_0 \otimes \dots \otimes a_{n-1}) = \begin{cases} a_0 \otimes \dots \otimes a_{i-1} \otimes \Delta(a_i) \otimes a_{i+1} \otimes \dots \otimes a_{n-1} \\ \text{if } i < n, \\ \tau(\Delta(a_0) \otimes a_1 \otimes \dots \otimes a_{n-1}) \\ \text{if } i = n, \end{cases}$$

where τ is the cyclic permutation toward the left, i.e.,

$$\tau(a_0 \otimes \dots \otimes a_n) = a_1 \otimes \dots \otimes a_n \otimes a_0.$$

The codegeneracy maps are constructed from the counit (augmentation) map on A in the usual way.

By the cyclic cobar complex of A, *denoted $coCH^\bullet(A, A)$, we mean the alternating sign cochain complex of the cyclic cobar construction on A. We write $coHH^*(A, A)$ for its cohomology, which we call* dual Hochschild cohomology.

Remark 2.3. Clearly, if B is a finite-dimensional co-commutative Hopf algebra over a field k, and if A is the k-linear dual of B, then the cyclic cobar complex on A is isomorphic to the k-linear dual of the cyclic bar complex on A, and the cyclic cobar complex then inherits a product from the coproduct on the cyclic bar complex of B.

It is worthwhile to be very explicit about the coproduct on the cyclic bar complex of a co-commutative Hopf algebra, since this point is not as well documented in the literature as it could be (although it does appear in some places, for example, in [1]). A Hochschild n-chain is an element of $A \otimes_k A \otimes_k \dots \otimes_k A$, with $n+1$ tensor factors of A; because the tensor factor on the far left plays a special role, we will adopt the common notation $a_0 [a_1 \otimes \dots \otimes a_n]$ instead of $a_0 \otimes a_1 \otimes \dots \otimes a_n$ for a (term in a) Hochschild n-chain.

Then the coproduct on the cyclic bar complex is given by

$$\Delta(a_0 [a_1 \otimes \dots \otimes a_n]) = \sum_{i=1}^{\ell} \sum_{j=0}^{n} a'_{0,i} \left[a_1 \otimes \dots \otimes a_j\right] \otimes a''_{0,i} \left[a_{j+1} \otimes \dots \otimes a_n\right]$$

where ℓ and $\{a'_{0,i}, a''_{0,i}\}_{i=1,\dots,\ell}$ are given by the determined by the coproduct in A and the formula

$$\Delta(a_0) = \sum_{i=0}^{\ell} a'_{0,i} \otimes a''_{0,i}.$$

We now give a definition of dual Hochschild cohomology with coefficients in the base field k, rather than in the coalgebra A itself. Naturally, one could write down a definition of dual Hochschild cohomology with coefficients in any "A-bicomodule," in a way that is basically obvious once one has taken a glance at Definitions 2.2 and 2.4. For the purposes of this paper, however, we will only ever need coefficients in k and in A.

Definition 2.4. *Let k be a field and A a copointed coalgebra over k, i.e., a coalgebra over k equipped with a morphism of k-coalgebras $\eta : k \to A$. By the* cyclic cobar construction *on A with coefficients in k we mean the cosimplicial k-vector space*

$$k \rightrightarrows k \otimes_k A \;\substack{\rightarrow\\\rightarrow\\\rightarrow}\; k \otimes_k A \otimes_k A \;\substack{\rightarrow\\\rightarrow\\\rightarrow\\\rightarrow}\; \dots$$

with coface maps $d^0, d^1, \ldots, d^n : A^{\otimes_k n} \to A^{\otimes_k n+1}$ *given by*

$$d^i(a_0 \otimes \ldots \otimes a_{n-1}) = \begin{cases} \tilde{\eta}_R(a_0) \otimes a_1 \otimes \ldots \otimes a_{n-1} \\ \text{if } i = 0, \\ a_0 \otimes \ldots \otimes a_{i-1} \otimes \Delta(a_i) \otimes a_{i+1} \otimes \ldots \otimes a_{n-1} \\ \text{if } 0 < i < n, \\ \tau\left(\tilde{\eta}_L(a_0) \otimes a_1 \otimes \ldots \otimes a_{n-1}\right) \\ \text{if } i = n, \end{cases}$$

where $\tilde{\eta}_R : k \to k \otimes_k A$ *is* η *composed with the usual isomorphism* $A \xrightarrow{\cong} k \otimes_k A$ *sending* a *to* $1 \otimes a$*, where* $\tilde{\eta}_L : k \to A \otimes_k k$ *is* η *composed with the usual isomorphism* $A \xrightarrow{\cong} A \otimes_k k$ *sending* a *to* $a \otimes 1$*, and where* τ *is as in Definition 2.2. The codegeneracy maps are constructed from the counit (augmentation) map on* A *in the usual way.*

By the cyclic cobar complex of A with coefficients in k*, denoted* $coCH^\bullet(A, k)$*, we mean the alternating sign cochain complex of the cyclic cobar construction on* A *with coefficients in* k*. We write* $coHH^*(A, k)$ *for its cohomology, which we call* dual Hochschild cohomology with coefficients in k.

Proposition 2.5. *Let* k *be a field and let* A *be a* k*-algebra which is finite-dimensional as a* k*-vector space. Let* A^* *denote the* k*-linear dual coalgebra of* A*. Then, for each* $n \in \mathbb{N}$*, the* n*th Hochschild homology* k*-vector space of* A *and the* n*th dual Hochschild cohomology* k*-vector space of* A^* *are mutually* k*-linearly dual. That is, we have isomorphisms of* k*-vector spaces:*

$$\hom_k(HH_n(A, A), k) \cong coHH^n(A^*, A^*), \tag{2.0.3}$$

$$\hom_k(coHH^n(A^*, A^*), k) \cong HH_n(A, A), \tag{2.0.4}$$

as well as isomorphisms

$$\hom_k(HH_n(A, k), k) \cong coHH^n(A^*, k), \tag{2.0.5}$$

$$\hom_k(coHH^n(A^*, k), k) \cong HH_n(A, k). \tag{2.0.6}$$

Proof. By construction, the cyclic cobar construction is simply the k-linear dual of the usual cyclic bar construction, so by the universal coefficient theorem (in its form for chain complexes), we have isomorphisms

$$\begin{aligned} coHH^n(A^*, A^*) &\cong H^n\left(coCH^\bullet(A, A)\right) \\ &\cong H^n\left(\hom_k(CH_\bullet(A, A), k)\right) \\ &\cong \hom_k\left(H_n(CH_\bullet(A, A)), k\right) \\ &\cong \hom_k\left(HH_n(A^*, A^*), k\right), \end{aligned}$$

giving us isomorphism 2.0.3. Finite-dimensionality of A as a k-vector space implies that $CH_n(A, A)$ is finite-dimensional as a k-vector space, hence the double dual of $CH_n(A, A)$ recovers $CH_n(A, A)$ again, giving us isomorphism 2.0.4. Essentially the same argument gives isomorphisms 2.0.5 and 2.0.6. □

Proposition 2.6. (May spectral sequence for dual Hochschild cohomology.) *Let* k *be a field and let* A *be a* k*-coalgebra. Let*

$$F_0A \subseteq F_1A \subseteq F_2A \subseteq \ldots \tag{2.0.7}$$

be a filtration of A which is comultiplicative, that is, if $x \in F_m A$, then $\Delta(x) \in \sum_{n=0}^{m} F_n A \otimes F_{m-n} A$. Then there exists a spectral sequence

$$E_1^{s,t} \cong coHH^{s,t}(E^0 A, E^0 A) \Rightarrow coHH^s(A,A)$$
$$d_r^{s,t} : E_r^{s,t} \to E_r^{s+1,t-r}.$$

The bigrading superscripts $coHH^{s,t}$ are as follows: s is the usual cohomological degree, while t is the May degree, defined and computed as follows: given a cohomology class $x \in coHH^s(E^0 A, E^0 A)$, its May degree is the total degree (in the grading on $E^0 A$ induced by the filtration on A) of any homogeneous cocycle representative for x in the cyclic cobar complex.

This spectral sequence enjoys the following additional properties:

(1) *If the filtration 2.0.7 is finite, i.e., $F_n A = 0$ for some $n \in \mathbb{N}$, then the spectral sequence converges strongly.*
(2) *If A is also a graded cocommutative k-coalgebra and the filtration layers $F_n A$ are generated (as two-sided coideals) by homogeneous elements, then this spectral sequence is a spectral sequence of graded k-vector spaces, i.e., the differential preserves the grading.*
(3) *If A is the underlying coalgebra of the k-linear dual Hopf algebra of a commutative Hopf algebra B over k, and the filtration 2.0.7 is a filtration by Hopf ideals, then $E^0 A$ is is also a commutative Hopf algebra, and the E_1-term and the abutment of the spectral sequence each have a natural ring structure. Furthermore, the spectral sequence is multiplicative, i.e., the differentials in the spectral sequence obey the graded Leibniz rule, and the product in the spectral sequence converges to the product on the abutment, modulo exotic multiplicative extensions (this is the usual situation in spectral sequences of differential graded algebras).*
(4) *The differential in the spectral sequence is (like any other spectral sequence of a filtered cochain complex) computed on a class $x \in coHH^{*,*}(E^0 A, E^0 A)$ by computing a homogeneous cocycle representative y for x in the cyclic cobar complex for $E^0 A$, lifting y to a homogeneous cochain $\tilde{y}$ in the cyclic cobar complex for A, applying the cyclic cobar differential d to $\tilde{y}$, then taking the image of $d\tilde{y}$ in the cyclic cobar complex for $E^0 A$.*

Proof. This is, of course, all formally dual to Proposition 2.1. The only thing that needs some explanation is the ring structure. The underlying filtered DGA of this spectral sequence has a ring structure given by the composite

$$\begin{aligned} coCH^\bullet(A,A) \otimes_k coCH^\bullet(A,A) &\xrightarrow{\cong} (CH_\bullet(B,B))^* \otimes_k (CH_\bullet(B,B))^* \\ &\xrightarrow{\cong} (CH_\bullet(B,B) \otimes_k CH_\bullet(B,B))^* \\ &\xrightarrow{\simeq} (CH_\bullet(B \otimes_k B, B \otimes_k B))^* \\ &\xrightarrow{\Delta^*} CH_\bullet(B,B)^* \\ &\xrightarrow{\cong} coCH^\bullet(A,A), \end{aligned}$$

where the map marked $\simeq$ is the k-linear dual of the Alexander-Whitney map, the map marked Δ^* is the k-linear dual of $CH_\bullet$ applied to the comultiplication map on B (which is well-defined, since $CH_\bullet$ is functorial on k-algebra maps and since B is assumed cocommutative, so that its comultiplication is a k-algebra morphism). The rest is formal. □

Proposition 2.7. (May spectral sequence for dual Hochschild cohomology, with coefficients in the base field.) *Let k be a field and let A be a k-coalgebra. Suppose A is equipped with a comultiplicative filtration as in 2.0.7. Then there exists a spectral sequence*

$$E_1^{s,t} \cong coHH^{s,t}(E^0A, k) \Rightarrow coHH^s(A, k)$$
$$d_r^{s,t} : E_r^{s,t} \to E_r^{s+1,t-r}.$$

The bigrading superscripts $coHH^{s,t}$ are as in Proposition 2.6.

This spectral sequence enjoys properties 1, 2, and 4 from Proposition 2.6.

Proof. Essentially identical to Proposition 2.6. □

3. The May and abelianizing filtrations.

We aim to compute $HH_*(A(1), A(1))$, the Hochschild homology of $A(1)$. By Proposition 2.5, this amounts to computing $coHH^*(A(1)^*, A(1)^*)$, and then taking the $\mathbb{F}_2$-linear dual. We now go about doing this.

Definition 3.1. *Recall that the* May filtration *on $A(1)$ is the filtration by powers of the augmentation ideal I. We write $\dot{F}^n(A(1))$ for the nth filtration layer in this filtration, i.e., $\dot{F}^nA(1) = I^n$, and we write $\dot{E}_0A(1)$ for the associated graded $\mathbb{F}_2$-algebra. If $x \in A(1)$, we sometimes write $\dot{x}$ for the associated element in $\dot{E}_0A(1)$.*

Proposition 3.2. *The $\mathbb{F}_2$-algebra $\dot{E}_0A(1)$ is the graded $\mathbb{F}_2$-algebra with generators $\dot{\mathrm{Sq}}^1$ and $\dot{\mathrm{Sq}}^2$ in grading degrees 1 and 2, respectively, and relations*

$$0 = \dot{\mathrm{Sq}}^1\dot{\mathrm{Sq}}^1 = \dot{\mathrm{Sq}}^2\dot{\mathrm{Sq}}^2 = \dot{\mathrm{Sq}}^1\dot{\mathrm{Sq}}^2\dot{\mathrm{Sq}}^1\dot{\mathrm{Sq}}^2 + \dot{\mathrm{Sq}}^2\dot{\mathrm{Sq}}^1\dot{\mathrm{Sq}}^2\dot{\mathrm{Sq}}^1.$$

The $\mathbb{F}_2$-linear duals of $A(1)$ and $\dot{E}_0(A(1))$ are, as Hopf algebras,

$$A(1)^* = \mathbb{F}_2[\overline{\xi}_1, \overline{\xi}_2]/\overline{\xi}_1^4, \overline{\xi}_2^2,$$
$$\Delta(\overline{\xi}_2) = \overline{\xi}_2 \otimes 1 + \overline{\xi}_1 \otimes \overline{\xi}_1^2 + 1 \otimes \overline{\xi}_2,$$
$$\left(\dot{E}_0A(1)\right)^* = \mathbb{F}_2[\overline{\xi}_{1,0}, \overline{\xi}_{1,1}, \overline{\xi}_{2,0}]/\overline{\xi}_{1,0}^2, \overline{\xi}_{1,1}^2, \overline{\xi}_{2,0}^2,$$
$$\Delta(\overline{\xi}_{2,0}) = \overline{\xi}_{2,0} \otimes 1 + \overline{\xi}_{1,0} \otimes \overline{\xi}_{1,1} + 1 \otimes \overline{\xi}_{2,0},$$

with $\overline{\xi}_1, \overline{\xi}_{1,0}, \overline{\xi}_{1,1}$ all primitive. The notation $\overline{x}$ is traditional for the conjugate of x in a Hopf algebra, and in this case, $\overline{\xi}_1 = \xi_1$ and $\overline{\xi}_2 = \xi_2 + \xi_1^3$. The notation $\overline{\xi}_{i,j}$ is used to denote the image of $\overline{\xi}_i^{2^j} \in A(1)$ in $\dot{E}_0(A(1))$.

Proof. Well-known consequence of the computation of the dual Steenrod algebra, as in [9]. □

Proposition 3.3. *The $\mathbb{F}_2$-algebra $\dot{E}_0A(1)$ is isomorphic to the group ring $\mathbb{F}_2[D_8]$ of the dihedral group D_8.*

Proof. We use the presentation

$$D_8 = \langle a, b \mid a^2, b^2, abab = baba \rangle$$

for D_8. The $\mathbb{F}_2$-algebra map

$$f : \mathbb{F}_2[D_8] \to \dot{E}_0A(1)$$

given by

$$f(a) = 1 + \dot{\mathrm{Sq}}^1$$
$$f(b) = 1 + \dot{\mathrm{Sq}}^2$$

is well-defined, since $f(a)^2 = 1 = f(b)^2$ and $f(a)f(b)f(a)f(b) = f(b)f(a)f(b)f(a)$. (Here it is essential that we are using $\dot{E}_0A(1)$ and not $A(1)$, since $\left(\dot{\mathrm{Sq}}^2\right)^2 = 0$ in $\dot{E}_0A(1)$ but $(\mathrm{Sq}^2)^2 \neq 0$ in $A(1)$.) One checks easily that $(a+1)^2 = 0 = (b+1)^2$ and $(a+1)(b+1)(a+1)(b+1) = (b+1)(a+1)(b+1)(a+1)$, hence we have a $\mathbb{F}_2$-algebra map

$$f^{-1} : \dot{E}_0A(1) \to \mathbb{F}_2[D_8]$$

sending $\dot{\mathrm{Sq}}^1$ to $a+1$ and sending $\dot{\mathrm{Sq}}^2$ to $b+1$. Clearly f^{-1} is inverse to f. □

We now use the well-known computation of the Hochschild homology of group rings (see e.g. Corollary 9.7.5 of [13]):

Theorem 3.4. *Suppose G is a discrete group, k a field. Let $\langle G\rangle$ be the set of conjugacy classes of elements in G, and given a conjugacy class S, let $C_G(S)$ denote the centralizer of S in G. Then there exists an isomorphism of graded k-vector spaces*

$$HH_*(k[G], k[G]) \cong \oplus_{S\in\langle G\rangle} H_*(C_G(S); k).$$

(Theorem 3.4 seems to be well-known, but I do not know who to attribute the result to, if anyone!)

Corollary 3.5. *The dimension of $HH_n(\dot{E}_0A(1), \dot{E}_0A(1))$ as a k-vector space is*

$$\dim_k HH_n(\dot{E}_0A(1), \dot{E}_0A(1)) = 3n + 5.$$

Proof. We use Proposition 3.3 and Theorem 3.4. There are five conjugacy classes of elements in $D_8 = \langle x, y \mid x^2, y^4, xy = y^3x\rangle$:

$$1, \{x, y^2x\}, \{yx, y^3x\}, \{y, y^3\}, \{y^2\},$$

with centralizers

$$D_8, \langle x, y^2 \mid x^2, (y^2)^2\rangle, \langle y^2 \mid (y^2)^2\rangle, \langle y \mid y^4\rangle, D_8,$$

respectively. These centralizer subgroups are isomorphic to

$$D_8, C_2 \times C_2, C_2, C_4, D_8,$$

respectively. The homology, with $\mathbb{F}_2$ coefficients, of these groups is well-known:

$$\dim_{\mathbb{F}_2} H_n(D_8; \mathbb{F}_2) = n + 1$$
$$\dim_{\mathbb{F}_2} H_n(C_2 \times C_2; \mathbb{F}_2) = n + 1$$
$$\dim_{\mathbb{F}_2} H_n(C_2; \mathbb{F}_2) = 1$$
$$\dim_{\mathbb{F}_2} H_n(C_4; \mathbb{F}_2) = 1,$$

hence

$$\begin{aligned}\dim_{\mathbb{F}_2} HH_n(\dot{E}_0A(1), \dot{E}_0A(1)) &= \dim_{\mathbb{F}_2} HH_n(\mathbb{F}_2[D_8], \mathbb{F}_2[D_8])\\ &= \dim_{\mathbb{F}_2} H_n(D_8; \mathbb{F}_2) + \dim_{\mathbb{F}_2} H_n(C_2 \times C_2; \mathbb{F}_2)\\ &+ \dim_{\mathbb{F}_2} H_n(C_2; \mathbb{F}_2) + \dim_{\mathbb{F}_2} H_n(C_4; \mathbb{F}_2)\\ &+ \dim_{\mathbb{F}_2} H_n(D_8; \mathbb{F}_2)\\ &= 3n + 5.\end{aligned}$$

□

Definition 3.6. *We now define a new filtration on $A(1)$ which we will call* the abelianizing filtration on $A(1)$. *To notationally distinguish it from the May filtration, we will write $\ddot{F}^n(A(1))$ for its filtration layers, and $\ddot{E}_0(A(1))$ for its associated graded algebra. The abelianizing filtration is defined as follows:*

$$\begin{aligned}
\ddot{F}^0(A(1)) &= A(1)\\
\ddot{F}^1(A(1)) &= (\mathrm{Sq}^1, \mathrm{Sq}^2)\\
\ddot{F}^2(A(1)) &= (\mathrm{Sq}^2)\\
\ddot{F}^3(A(1)) &= (\mathrm{Sq}^1\,\mathrm{Sq}^2, \mathrm{Sq}^2\,\mathrm{Sq}^1)\\
\ddot{F}^4(A(1)) &= (Q_0)\\
\ddot{F}^5(A(1)) &= (\mathrm{Sq}^1\,Q_0, \mathrm{Sq}^2\,Q_0)\\
\ddot{F}^6(A(1)) &= (\mathrm{Sq}^2\,Q_0)\\
\ddot{F}^7(A(1)) &= (\mathrm{Sq}^1\,\mathrm{Sq}^2\,Q_0)\\
\ddot{F}^8(A(1)) &= 0,
\end{aligned}$$

where (x) is two-sided ideal generated by an element x, and where Q_0 is Milnor's notation for the element $\mathrm{Sq}^1\,\mathrm{Sq}^2 + \mathrm{Sq}^2\,\mathrm{Sq}^1$. If $x \in A(1)$, we sometimes write $\ddot{x}$ for the associated element in $\ddot{E}_0A(1)$.

Proposition 3.7. *The abelianizing filtration on $A(1)$ has the following properties:*

- *The abelianizing filtration is finer than the May filtration, that is, $\dot{F}^n(A(1)) \subseteq \ddot{F}^n(A(1))$ for all n.*
- *The abelianizing filtration is a multiplicative filtration, that is, if $x \in \ddot{F}^m(A(1))$ and $y \in \ddot{F}^n(A(1))$, then $xy \in \ddot{F}^{m+n}(A(1))$.*
- *Furthermore, the abelianizing filtration is a Hopf filtration, that is, if $x \in \ddot{F}^m(A(1))$, then*
$$\Delta(x) \in \sum_{i=0}^{m} \ddot{F}^i(A(1)) \otimes_{\mathbb{F}_2} \ddot{F}^{m-i}(A(1)).$$
- *The associated graded Hopf algebra $\ddot{E}_0(A(1))$ of the abelianizing filtration on $A(1)$ is the exterior algebra $E(\ddot{\mathrm{Sq}}^1, \ddot{\mathrm{Sq}}^2, \ddot{Q}_0)$, with $\ddot{\mathrm{Sq}}^1$ and $\ddot{Q}_0$ primitive, and with $\Delta(\ddot{\mathrm{Sq}}^2) = \ddot{\mathrm{Sq}}^2 \otimes 1 + \ddot{\mathrm{Sq}}^1 \otimes \ddot{\mathrm{Sq}}^1 + 1 \otimes \ddot{\mathrm{Sq}}^2$. The topological degrees (i.e., inherited from the usual grading on the Steenrod algebra) of $\ddot{\mathrm{Sq}}^1, \ddot{\mathrm{Sq}}^2, \ddot{Q}_0$ are $1, 3$, and 4, respectively, and their abelianizing degrees (i.e., inherited from the abelianizing filtration) are $1, 2$, and 4, respectively.*
- *The $\mathbb{F}_2$-linear dual Hopf algebra $\left(\ddot{E}_0(A(1))\right)^*$ is*
$$\left(\ddot{E}_0(A(1))\right)^* \cong \mathbb{F}_2[\overline{\ddot{\xi}}_{1,0}, \overline{\ddot{\xi}}_{2,0}]/\overline{\ddot{\xi}}_{1,0}^4, \overline{\ddot{\xi}}_{2,0}^2,$$
with $\overline{\ddot{\xi}}_{1,0}, \overline{\ddot{\xi}}_{2,0}$ both primitive. In particular, $\left(\ddot{E}_0(A(1))\right)^*$ is isomorphic to $\left(\dot{E}_0(A(1))\right)^*$ as $\mathbb{F}_2$-coalgebras (but not as Hopf algebras).

Proof. Elementary computation. □

4. Running the HH-May and abelianizing spectral sequences.

4.1. **Input.**

Lemma 4.1. *Let k be a field of characteristic two, and give $k[x]/x^2$ the structure of a Hopf algebra over k by letting $\Delta(x) = x \otimes 1 + 1 \otimes x$. Then we have an isomorphism of graded Hopf algebras*

$$HH_*(k[x]/x^2, k[x]/x^2) \cong k[x]/x^2 \otimes_k \Gamma_k(\sigma x),$$

where:

- *$HH_*(k[x]/x^2, k[x]/x^2)$ inherits its coproduct from that of $k[x]/x^2$, as in Remark 2.3,*
- *$\Gamma_k(\sigma x)$ denotes the divided power algebra over k on a generator σx,*
- *σx denotes the homology class of the 1-cycle $1[x]$, i.e., $1 \otimes x$ (and consequently σx is in homological degree 1),*
- *$\Delta(x) = x \otimes 1 + 1 \otimes x$, and*
- *$\Delta(\gamma_n(\sigma x)) = \sum_{i=0}^{n} \gamma_i(\sigma x) \otimes \gamma_{n-i}(\sigma x)$, where γ_n is the nth divided power.*

Consequently, the linear dual of $HH_*(k[x]/x^2, k[x]/x^2)$, i.e., the cohomology of the cyclic cobar complex of the linear dual of $k[x]/x^2$, is isomorphic as a Hopf algebra to $k[\xi]/\xi^2 \otimes_k k[h]$, with ξ, h both primitive, where ξ is the dual basis element to x, and h is the dual basis element to σx; consequently ξ, h are in grading degrees $0, 1$, respectively.

Proof. Using the cycle representative $1\,[x \otimes \ldots \otimes x]$ for $\gamma_n(\sigma x)$, with n tensor factors of x, the claims made are easy (and classical) computations of coproducts (using Remark 2.3) and shuffle products in the cyclic bar complex. □

Lemma 4.2. *Let k be a field of characteristic two, and give $k[x,y]/(x^2,y^2)$ the structure of a Hopf algebra over k by letting $\Delta(x) = x \otimes 1 + 1 \otimes x$ and by letting $\Delta(y) = y \otimes 1 + x \otimes x + 1 \otimes y$. Then we have an isomorphism of graded Hopf algebras*

$$HH_*\left(k[x,y]/(x^2,y^2), k[x,y]/(x^2,y^2)\right) \cong k[x,y]/(x^2,y^2) \otimes_k \Gamma_k(\sigma x, \sigma y),$$

where:

- *$HH_*(k[x,y]/(x^2,y^2), k[x,y]/(x^2,y^2))$ inherits its coproduct from that of $k[x,y]/(x^2,y^2)$, as in Remark 2.3,*
- *$\Gamma_k(\sigma x, \sigma y)$ denotes the tensor product over k of the divided power algebra over k on a generator σx with the divided power algebra over k on a generator σy,*
- *σx (respectively, σy) denotes the homology class of the 1-cycle $1[x]$ (respectively, $1[y]$), and consequently σx and σy are each in homological degree 1,*
- *the coproducts $\Delta(x)$ and $\Delta(y)$ are exactly as in $k[x,y]/(x^2,y^2)$,*
- *$\Delta(\gamma_n(\sigma x)) = \sum_{i=0}^{n} \gamma_i(\sigma x) \otimes \gamma_{n-i}(\sigma x)$, and*
- *$\Delta(\gamma_n(\sigma y)) = \sum_{i=0}^{n} \gamma_i(\sigma y) \otimes \gamma_{n-i}(\sigma y)$.*

Consequently, the linear dual of $HH_*(k[x,y]/(x^2,y^2), k[x,y]/(x^2,y^2))$, i.e., the cohomology of the cyclic cobar complex of the linear dual of $k[x,y]/(x^2,y^2)$, is isomorphic as a Hopf algebra to $k[\xi]/\xi^4 \otimes_k k[h_x, h_y]$, with ξ, h_x, h_y all primitive, where ξ is the dual basis element to x, and h_x, h_y are the dual basis elements to $\sigma x, \sigma y$, respectively; consequently ξ, h_x, h_y are in grading degrees $0, 1, 1$, respectively.

Proof. Just as in Lemma 4.1. (Note that $\Delta(\gamma_n(\sigma y))$ does not involve x or σx, even though $\Delta(y)$ involves x; this is simply because the formula for $\Delta(\sigma y) = \Delta\,(1\,[y])$, given in Remark 2.3, does not actually make any use of the coproduct of y in $k[x,y]/(x^2,y^2)$.) □

Proposition 4.3. *There exist four strongly convergent trigraded multiplicative spectral sequences:*

(Abelianizing:)

$$E_1^{s,t,u} \cong coHH^{s,t,u}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right) \Rightarrow coHH^s(A(1)^*, A(1)^*)$$
$$E_1^{*,*,*} \cong \mathbb{F}_2[x_{10}]/(x_{10}^4) \otimes_{\mathbb{F}_2} E(x_{20}) \otimes_{\mathbb{F}_2} P(h_{10}, h_{11}, h_{20})$$
$$d_r^{s,t,u} : E_r^{s,t,u} \to E_r^{s+1,t-r,u},$$

(HH-May:)

$$E_1^{s,t,u} \cong coHH^{s,t,u}\left(\dot{E}^0(A(1)^*), \dot{E}^0(A(1)^*)\right) \Rightarrow coHH^s(A(1)^*, A(1)^*)$$
$$d_r^{s,t,u} : E_r^{s,t,u} \to E_r^{s+1,t-r,u},$$

(HH-May with coeffs. in $\mathbb{F}_2$:)

$$E_1^{s,t,u} \cong coHH^{s,t,u}\left(\dot{E}^0(A(1)^*), \mathbb{F}_2)\right) \Rightarrow coHH^s(A(1)^*, \mathbb{F}_2)$$
$$d_r^{s,t,u} : E_r^{s,t,u} \to E_r^{s+1,t-r,u},$$

(Abelianizing-to-HH-May:)

$$E_1^{s,t,u} \cong coHH^{s,t,u}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right) \Rightarrow coHH^s(\dot{E}^0A(1)^*, \dot{E}^0A(1)^*)$$
$$E_1^{*,*,*} \cong \mathbb{F}_2[x_{10}]/(x_{10}^4) \otimes_{\mathbb{F}_2} E(x_{20}) \otimes_{\mathbb{F}_2} P(h_{10}, h_{11}, h_{20})$$
$$d_r^{s,t,u} : E_r^{s,t,u} \to E_r^{s+1,t-r,u},$$

where P denotes a polynomial algebra (over $\mathbb{F}_2$) and E an exterior algebra (over $\mathbb{F}_2$), and there exists a morphism of spectral sequences from the HH-May spectral sequence to the HH-May spectral sequence with coefficients in $\mathbb{F}_2$. We write s for the cohomological degree, t for the filtration degree, and u for the internal/topological degree. Here (and elsewhere) we write x_{i0} for the cohomology class of the cyclic cobar 0-cocycle $\overline{\xi}_{i,0}$ and h_{ij} for the cohomology class of cyclic cobar 1-cocycle $1\left[\overline{\xi}_{i,j}\right]$, where $\overline{\xi}_{1,0}, \overline{\xi}_{1,1}, \overline{\xi}_{2,0}$ are the dual basis elements to the elements $\ddot{\mathrm{Sq}}^1, \ddot{\mathrm{Sq}}^2, \ddot{Q}_0$, respectively, in $\ddot{E}_0A(1)$. (Of course the machinery we are using produces much more than just these four spectral sequences; but these four are the ones we will actually use in this paper.)

The tridegrees of the generators of $coHH^{*,*,*}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right)$, and cocycle representatives for those cohomology classes in the cyclic cobar complex, are as follows:

$$x_{10} = [\overline{\xi}_{1,0}] \in coCH^{0,1,1}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right)$$
$$x_{20} = [\overline{\xi}_{2,0}] \in coCH^{0,4,3}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right)$$
$$h_{10} = [1 \otimes \overline{\xi}_{1,0}] \in coCH^{1,1,1}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right)$$
$$h_{11} = [1 \otimes \overline{\xi}_{1,1}] \in coCH^{1,2,2}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right)$$
$$h_{20} = [1 \otimes \overline{\xi}_{2,0}] \in coCH^{1,4,3}\left(\ddot{E}^0(A(1)^*), \ddot{E}^0(A(1)^*)\right)$$

Proof. Consequence of Propositions 2.6, 3.2 and 3.7 and Lemmas 4.1 and 4.2. □

Proposition 4.4. *The HH-May spectral sequence with coefficients in $\mathbb{F}_2$, from Proposition 4.3, is isomorphic (beginning with the E_1-term) to the classical May spectral sequence for $A(1)$, $\mathrm{Ext}^{*,*,*}_{E^0A(1)}(\mathbb{F}_2,\mathbb{F}_2) \Rightarrow \mathrm{Ext}^{*,*}_{A(1)}(\mathbb{F}_2,\mathbb{F}_2)$. (See e.g. Example 3.2.7 of* [11] *for this spectral sequence.)*

Proof. The cyclic cobar complex of $A(1)$ with coefficients in $\mathbb{F}_2$ is isomorphic to the classical (non-cyclic) cobar complex of $A(1)$, as in Definition A.1.2.11 of [11]. and the May filtration on one coincides with the May filtration on the other. □

4.2. d_1**-differentials.**

Proposition 4.5. *In both the abelianizing spectral sequence and the abelianizing-to-HH-May spectral sequence, the d_1 differentials are given on the multiplicative generators by*

$$\begin{aligned}
d_1(x_{10}) &= 0,\\
d_1(x_{20}) &= x_{10}h_{11} + x_{10}^2 h_{10},\\
d_1(h_{10}) &= 0,\\
d_1(h_{11}) &= 0, \text{ and}\\
d_1(h_{20}) &= h_{10}h_{11}.
\end{aligned}$$

Using these formulas and the Leibniz rule, we get the d_1 differential on all elements of the E_1-terms of the abelianizing and abelianizing-to-HH-May spectral sequences.

Proof. In Proposition 4.3 we gave cocycle representatives for the six multiplicative generators. We then easily compute the d_1 differentials on those generators using the method described in Proposition 2.6:

$$\begin{aligned}
d(\overline{\xi}_{1,0}) &= 0,\\
d(\overline{\xi}_{2,0}) &= \overline{\xi}_{1,0}\otimes\overline{\xi}_{1,1} + \overline{\xi}_{1,1}\otimes\overline{\xi}_{1,0}\\
&\quad + 1\otimes\overline{\xi}_{2,0} + \overline{\xi}_{2,0}\otimes 1\\
&\quad + \overline{\xi}_{2,0}\otimes 1 + 1\otimes\overline{\xi}_{2,0}\\
&= \overline{\xi}_{1,0}\otimes\overline{\xi}_{1,1} + \overline{\xi}_{1,1}\otimes\overline{\xi}_{1,0}\\
d(1\otimes\overline{\xi}_{1,0}) &= 0,\\
d(1\otimes\overline{\xi}_{1,1}) &= 0,\\
d(1\otimes\overline{\xi}_{2,0}) &= 1\otimes 1\otimes\overline{\xi}_{2,0} + 1\otimes\overline{\xi}_{2,0}\otimes 1\\
&\quad + 1\otimes\overline{\xi}_{1,0}\otimes\overline{\xi}_{1,1} + 1\otimes 1\otimes\overline{\xi}_{2,0}\\
&\quad + 1\otimes\overline{\xi}_{2,0}\otimes 1\\
&= 1\otimes\overline{\xi}_{1,0}\otimes\overline{\xi}_{1,1}.
\end{aligned}$$

Here $\overline{\xi}_{1,1}$ is the dual basis element to Sq^2, so $\overline{\xi}_{1,1} = \overline{\xi}_{1,0}^2$ in the associated graded of the abelianizing filtration, but $\overline{\xi}_{1,1}$ is indecomposable in the associated graded of the May filtration. Using the product on dual Hochschild cohomology from Proposition 2.6, we get that these cocycles represent the cohomology classes $0, 0, x_{10}h_{11}+x_{10}^2h_{10}, 0, h_{10}h_{11}$, respectively. □

Now one has enough information to do a routine computation of the cohomology of the E_1-term, and get the E_2-term. While h_{20} is not a cocycle in the E_1-term, its square is,

and we follow the traditional (due to May's thesis) notational conventions of May spectral sequences by writing b_{20} for h_{20}^2.

We present the E_2-term as a spectral sequence chart.

Conventions 4.6. In all the spectral sequence charts in this paper,

- the vertical axis is the homological degree,
- the horizontal axis is the Adams degree, i.e., the internal/topological degree minus the homological degree,
- horizontal lines (whether curved or straight) represent multiplication by x_{10},
- vertical lines represent multiplication by h_{10}, and
- diagonal lines represent multiplication by h_{11}.

Here is a spectral sequence chart illustrating the E_2-term of the abelianizing and abelianizing-to-HH-May spectral sequences (their E_2-terms are abstractly isomorphic as bigraded $\mathbb{F}_2$-vector spaces, if one forgets about the filtration degree and only keeps track of the cohomological and Adams degrees), reduced modulo the ideal generated by b_{20}:

(4.2.1)



The classes whose names are not implied by the lines representing various multiplications are as follows:

- the class in bidegree $(1, 5)$ is $x_{10}^2(h_{10}x_{20} + x_{10}h_{20})$, which we abbreviate as z,
- and the class in bidegree $(0, 6)$ is $x_{10}^3 x_{20}$, which we abbreviate as x_6.

The spectral sequence's E_2-term is b_{20}-periodic, that is, there exists a class (not pictured) b_{20} in bidegree $(2, 4)$ each of whose positive integer powers generates an isomorphic copy of the chart 4.2.1.

Consequently, as a trigraded $\mathbb{F}_2$-algebra, the spectral sequence's E_2-term is isomorphic to:

$$\begin{aligned}\mathbb{F}_2[x_{10}, h_{10}, h_{11}, z, x_6, b_{20}] \text{ modulo relations } & x_{10}^4,\ x_{10}h_{11} = x_{10}^2 h_{10},\\ & h_{10}h_{11},\ x_{10}z = h_{10}x_6,\ h_{11}z,\ z^2,\\ & x_{10}x_6,\ zx_6,\ x_6^2,\end{aligned}$$

with generators in tridegrees:

Class	Coh. degree	Ab. degree	Top. degree	Adams degree
x_{10}	0	1	1	1
x_6	0	7	6	6
h_{10}	1	1	1	0
h_{11}	1	2	2	1
z	1	7	6	5
b_{20}	2	8	6	4

4.3. d_2**-differentials.**

Proposition 4.7. *The abelianizing-to-HH-May spectral sequence collapses at E_2, i.e., there are no nonzero differentials longer than d_1 differentials. Consequently, the spectral sequence chart 4.2.1 describes the E_1-term (and also the E_2-term) of the HH-May spectral sequence, as well as the E_2-term of the abelianizing spectral sequence.*

Proof. An easy dimension count on the E_2-term 4.2.1 gives us that the $\mathbb{F}_2$-vector space dimension of the s-row is $3s + 5$. By Proposition 2.5 and Corollary 3.5, this is the correct dimension for the E_∞-term. So there can no further nonzero differentials in the spectral sequence, since any such differentials would reduce the $\mathbb{F}_2$-vector space dimension of some row. □

Proposition 4.8. *The d_2 differentials on the multiplicative generators of the E_2-term of the HH-May spectral sequence, as well as the abelianizing spectral sequence, are as follows:*

$$\begin{aligned} d_2(x_{10}) &= 0, \\ d_2(h_{10}) &= 0, \\ d_2(h_{11}) &= 0, \\ d_2(z) &= 0, \\ d_2(x_6) &= 0, \text{ and} \\ d_2(b_{20}) &= h_{11}^3. \end{aligned}$$

Using these formulas and the Leibniz rule, we get the d_2 differential on all elements of the E_2-term of the abelianizing spectral sequence.

Proof. For x_{10}, h_{10}, and h_{11}, same computation as Proposition 4.5. For z, x_6, inspection of the tridegrees of elements rules out all nonzero possibilities for d_2.

For the differential $d_2(b_{20})$: we see from inspection of the tridegrees that the only possible nonzero differential on b_{20} would have to hit a scalar multiple of h_{11}^3, and this differential indeed occurs, using Proposition 4.4 to map the HH-May spectral sequence to the classical May spectral sequence for $A(1)$, in which the differential $d_2(b_{20}) = h_{11}^3$ is classical and well-known (see e.g. Lemma 3.2.10 of [11]). □

So the only nonzero d_2 differentials are the d_2-differential $d_2(b_{20}) = h_{11}^3$ and its products with other classes. By the Leibniz rule, $d_2(b_{20}^2) = 0$, so the spectral sequence's E_3-term is b_{20}^2-periodic. We now draw a chart illustrating the E_3-term, modulo the two-sided ideal generated by b_{20}^2:

(4.3.1)



The entire pattern described by the chart 4.3.1 repeats: there is the periodicity class (not pictured) b_{20}^2 in bidegree $(4,8)$, which maps, under the map of spectral sequences of Proposition 4.4, to the element in $\mathrm{Ext}_{A(1)}^{4,8}(\mathbb{F}_2,\mathbb{F}_2)$ which is the image in the associated graded of the Adams filtration of the famous real Bott periodicity element in $\pi_8(ko)$.

The classes whose names are not implied by the lines representing various multiplications are as follows, and whose names were not already given in our description of the E_2-term, are as follows:

- the class in bidegree $(4,3)$ is $h_{10}b_{20}$, which we abbreviate as w_4,
- the class in bidegree $(5,2)$ is $x_{10}b_{20}$, which we abbreviate as w_5,
- and the class in bidegree $(9,3)$ is zb_{20}, which we abbreviate as w_9.

Finally, we write b for b_{20}^2, so that the spectral sequence's E_3-term is multiplicatively generated by elements:

Class	Coh. degree	Ab. degree	Top. degree	Adams degree
x_{10}	0	1	1	1
x_6	0	7	6	6
h_{10}	1	1	1	0
h_{11}	1	2	2	1
z	1	7	6	5
w_4	3	9	7	4
w_5	2	9	7	5
w_9	3	15	12	9
b	4	16	12	8 .

In Proposition 4.4 we constructed a map from the HH-May spectral sequence to the classical May spectral sequence computing $\mathrm{Ext}_{A(1)}^{*,*}(\mathbb{F}_2,\mathbb{F}_2)$. We now draw the $E_3 \cong E_\infty$-term of that classical May spectral sequence, using the same conventions as charts 4.2.1 and 4.3.1, so that one can easily see the (surjective) map of spectral sequence E_3-terms:

(4.3.2)



Again, there is a periodicity class (not pictured) $b = b_{20}$ in bidegree $(4, 8)$, i.e., cohomological degree 4 and topological degree 12 (hence Adams degree 8).

Proposition 4.9. *In the abelianizing and the HH-May spectral sequences, all d_r differentials are zero, for all $r > 2$.*

Proof. We simply check that there can no nonzero d_r differentials, for $r > 2$, on the multiplicative generators $x_{10}, x_6, h_{10}, h_{11}, z, w_4, w_5, w_9, b$ of the E_3-term of the abelianizing, equivalently (starting with E_3), the HH-May spectral sequence. In the proof of Proposition 4.8, we showed that x_{10}, h_{10}, h_{11}, and x_6 all do not support differentials of any length whatsoever. The remaining classes are all incapable of supporting nonzero d_r differentials, for $r > 2$, for degree reasons: there are no classes in the correct tridegree for any of these classes to hit by a d_r differential, if $r > 2$. □

Theorem 4.10. *The spectral sequence chart 4.3.1 displays (by reading across the rows) the Hochschild homology $HH_*(A(1), A(1))$. In particular, the $\mathbb{F}_2$-vector space dimension of $HH_n(A(1), A(1))$ is:*

$$\dim_{\mathbb{F}_2} HH_n(A(1), A(1)) = \begin{cases} 2n+5 & \text{if } 2 \mid n \\ 2n+6 & \text{if } n \equiv 1 \mod 4 \\ 2n+4 & \text{if } n \equiv 3 \mod 4. \end{cases}$$

Hence the Poincaré series of the graded $\mathbb{F}_2$-vector space $HH_*(A(1), A(1))$ is

$$\frac{5 + 8s + 9s^2 + 10s^3 + \frac{8s^4}{1-s}}{1 - s^4}.$$

If we additionally keep track of the extra grading on $HH_*(A(1), A(1))$ coming from the topological grading on $A(1)$, then the Poincaré series of the bigraded $\mathbb{F}_2$-vector space $HH_{*,*}(A(1), A(1))$ is

$$\left((1+u)\left(1+u^2+su(1+u^2+u^5)+s^2u^2(1+2u^5+u^7)+s^3u^3(1+u^4+u^5+u^6+u^9) + \frac{s^4u^4(1+u^4+u^5+u^9)}{1-su}\right) + u^6 + su^2 + su^8 + s^2u^4\right)\frac{1}{1-s^4u^{12}} \tag{4.3.3}$$

where s indexes the homological grading and u indexes the topological grading, as in Proposition 4.3.

Proof. This information is read off directly from the spectral sequence chart 4.3.1. (Note that the horizontal axis in the chart 4.3.1 is the Adams degree, i.e., $u - s$, not the internal/topological degree, i.e., u, so one must be a little careful in reading off the series 4.3.3 from the chart.) □

References

[1] Hossein Abbaspour. On the Hochschild homology of open Frobenius algebras. *available on arXiv*.

[2] Jean-Luc Brylinski. A differential complex for Poisson manifolds. *J. Differential Geom.*, 28(1):93–114, 1988.

[3] Bjørn Ian Dundas, Thomas G. Goodwillie, and Randy McCarthy. *The local structure of algebraic K-theory*, volume 18 of *Algebra and Applications*. Springer-Verlag London, Ltd., London, 2013.

[4] S. Geller, L. Reid, and C. Weibel. The cyclic homology and K-theory of curves. *J. Reine Angew. Math.*, 393:39–90, 1989.

[5] Ib Madsen. Algebraic K-theory and traces. In *Current developments in mathematics, 1995 (Cambridge, MA)*, pages 191–321. Int. Press, Cambridge, MA, 1994.

[6] W. S. Massey. Products in exact couples. *Ann. of Math. (2)*, 59:558–569, 1954.

[7] J. Peter May. *The cohomology of restricted Lie algebras and of Hopf algebras: application to the Steenrod algebra*. ProQuest LLC, Ann Arbor, MI, 1964. Thesis (Ph.D.)–Princeton University.

[8] J. E. McClure and R. E. Staffeldt. On the topological Hochschild homology of bu. I. *Amer. J. Math.*, 115(1):1–45, 1993.

[9] John Milnor. The Steenrod algebra and its dual. *Ann. of Math. (2)*, 67:150–171, 1958.

[10] Teimuraz Pirashvili and Friedhelm Waldhausen. Mac Lane homology and topological Hochschild homology. *J. Pure Appl. Algebra*, 82(1):81–98, 1992.

[11] Douglas C. Ravenel. *Complex cobordism and stable homotopy groups of spheres*, volume 121 of *Pure and Applied Mathematics*. Academic Press Inc., Orlando, FL, 1986.

[12] N. E. Steenrod. *Cohomology operations*. Lectures by N. E. Steenrod written and revised by D. B. A. Epstein. Annals of Mathematics Studies, No. 50. Princeton University Press, Princeton, N.J., 1962.

[13] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.

[14] Charles A. Weibel. *The K-book*, volume 145 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K-theory.